



Vail, C. S., Reiss, D., Brakes, P., & Butterworth, A. (2019). Potential Welfare Impacts of Chase and Capture of Small Cetaceans during Drive Hunts in Japan. *Journal of Applied Animal Welfare Science*.
<https://doi.org/10.1080/10888705.2019.1574576>

Peer reviewed version

Link to published version (if available):
[10.1080/10888705.2019.1574576](https://doi.org/10.1080/10888705.2019.1574576)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Taylor and Francis at
<https://www.tandfonline.com/doi/abs/10.1080/10888705.2019.1574576?needAccess=true>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Potential Welfare Impacts of Chase and Capture of Small Cetaceans During Drive Hunts in Japan

Courtney S. Vail¹, Diana Reiss², Philippa Brakes^{3, 4}, Andrew Butterworth⁵

¹Lightkeepers, Phoenix, AZ, USA; ²Department of Psychology, Hunter College, City University of New York, New York, NY, USA; ³Centre for Ecology and Conservation, University of Exeter, Tremough, Cornwall, UK; ⁴Whale and Dolphin Conservation (WDC), Wiltshire, UK; ⁵Clinical Veterinary School, University of Bristol Veterinary School, Langford, UK

Abstract

Drive hunts are a method to herd, capture and kill small cetaceans (whales and dolphins) in coastal waters of some countries including Japan and the Faroe Islands. In Japan, these methods are often associated with the acquisition of live dolphins for international marine parks and aquaria. During the hunts, dolphins are herded by a flotilla of fishing vessels and loud underwater noise created by fishermen banging hammers on metal poles. The prolonged and strenuous chase and use of sound barriers to herd, capture, and restrain the dolphins can result in acute stress and injury. The authors review physiological and behavioural data pertaining to chase, encirclement, and live capture of dolphins and draw comparisons between chase and capture data for marine and terrestrial species. This analysis raises substantial welfare concerns associated with the hunts and acquisition of dolphins from such capture operations. The authors assert that this data detailing the negative impacts of chase, herding and handling (capture) of small cetaceans renders these hunts inherently inhumane and should inform policy relating to the collection and management of dolphins in the wild.

Introduction

Drive hunts are a method used to herd, capture and kill small cetaceans (whales and dolphins) in the coastal waters of some countries including Japan and the Faroe Islands. In Japan, the animals are herded by a flotilla of fishing vessels (Figure 1) and a cacophony of underwater noise is created by fishermen banging hammers on specifically designed trumpet-shaped metal poles (Figures 2 (a,b)). The practice generates powerful acoustic signals (greater than 170 dB) which are transmitted through the water inducing flight, escape and avoidance behavior in the animals pursued in these hunts. This is known as the Oikomi method (Akamatsu, Hatakeyama, & Takatsu, 1993; Brownell, Nowacek, & Ralls, 2008; Ohsumi, 1972) and it is used to drive cetaceans into harbors or shallow coves where they will be either killed, or selected for display in marine parks and aquaria. The main species taken in the Japanese drive hunts include: common bottlenose dolphins (*Tursiops truncatus*), striped dolphins (*Stenella coeruleoalba*), Risso's dolphins (*Grampus griseus*), false killer whales (*Pseudorca crassidens*), Pacific white-sided dolphins (*Lagenorhynchus obliquidens*), pantropical spotted dolphins (*Stenella attenuata*), and short-finned pilot whales (*Globicephala macrorhynchus*). More recently, melon-headed whales (*Peponocephala electra*) and rough-toothed dolphins (*Steno bredanensis*) have been added to the targeted species list. Quotas of over 2,000 dolphins and small whales are issued annually for these specific hunts (Kasuya, 2007).

In a previous report, we describe and discuss the killing methods and techniques employed in the drive hunts and why they do not conform to the recognized requirements for humane killing methods that render “immediate insensibility” and thus would not be permitted in any regulated slaughterhouse process in the developed world (Butterworth et al., 2013).



Figure 1. A fleet of boats returns from a dolphin drive hunt in Taiji, Japan. Image: DolphinProject.com



Figure 2. (a) Metal trumpets utilized in the dolphin drive hunts. (b) Trumpets (indicated by circle) are mounted on vessels and placed in the water. Using a hammer or other object, fishermen bang on the poles to create loud noise (up to 205dB) that disorients the dolphins. Image: Courtney Vail

In this paper, we focus on the chasing, herding, separation and confinement of animals during the drive hunt process, and review the existing literature and other available data on the physical, behavioral and social impacts of these activities on cetaceans. This data is compared to existing data on the impact of hunt and acquisition activities in other species. We raise questions about the potential welfare and conservation implications of the methods used in the drive hunts and highlight the notable absence of scientific data applied in policies governing dolphin management and acquisition.

Acoustic Impacts: an aversive wall of sound

During the drive hunts in Japan, the aversive wall of sound is created by the combination of boat engine noise and repetitive banging on the long metal trumpets.

Brownell et al. (2008, p.82) report that:

“When a scouting boat finds a school, it reports to the office on shore and 10 to 20 driving boats are sent to surround the school and drive it, by making noise underwater with ‘trumpets’ (Kasuya, 2002), into the bay of the village where the dolphins are stranded on the beach or surrounded with a long net and killed.”

Akamatsu et al. (1993) assessed the action of the metal trumpets and showed that the sound was effective in causing the animals to swim directly away from the source at received levels of 170 dB re: 1mPa. The same researcher also reported that these trumpets, also known as banger poles, could produce sound at 205dB re: 1mPa at 0.2-5.2kHz. Both 174dB and 205 dB are ‘extremely loud’ sound levels (louder than a jet engine at close proximity) and in shallow water and where the sound propagates outward in a concentric ring, cetaceans at ranges of up to approx. 1200m (1.2km) can potentially receive sound levels capable of eliciting strong behavioral responses. Brownell et al. (2008, p. 86) concluded that:

“It is not surprising that Japanese fishermen, using two of these [trumpets] on each of several vessels at the same time, are able to herd schools of small cetaceans from tens of kilometres offshore into harbours, or cause them to mass-strand on beaches.”

Anthropogenic sound is a potential stressor for marine mammals that affects psychological and physical health, as has been demonstrated in other mammals (Romano et al., 2002; Tyack, 2008). The primary sense of cetaceans is hearing and they are reliant on sound continuously for multiple social and biological functions such as foraging, navigation, maintaining social contact with members of their social group, breeding, and threat avoidance. The exposure to high levels of noise, and especially to prolonged exposure to noise, will negatively impact many of these important functions (Tyack & Clark, 2000; Wright et al., 2007a, 2007b). Noise, has been defined as “essentially as an aperiodic signal that interferes with the perception of sound and has a negative physiological impact” (Ketten, 2016, p.208).

Although cetaceans use a rich variety of multi-modal signals (e.g., behavioral, spatial, and tactile), they rely heavily on acoustic communication to maintain contact, both at close range, and also at longer distances. Many toothed cetaceans use wide-band sonar signals for navigation and orientation and use a rich repertoire of frequency-modulated whistles and pulsed calls for communication and cohesion between group members (Brown, Caldwell, & Caldwell, 1966; Esch, Sayigh, Blum, & Wells, 2009; McCowan & Reiss, 1995). With disrupted capacity to communicate, the animals may experience social disorientation, increased stress and loss of ability to use acoustic senses (Tyack, 2008; Weilgart, 2007). Elevated background noise levels have been identified as a source of interference in marine mammals’ ability to detect calls from other individuals, and may result in long-term displacement or abandonment of important habitat where resting, feeding, breeding, migratory patterns, and social interactions are interrupted through noise disturbance (Fair & Becker, 2000; Nowacek, 2007; Richardson & Wursig, 1997; Tyack, 2008; Weilgart, 2007).

It has been well-documented that repeated exposure to loud noises may result in physical hearing loss. Noise-induced ear damage has been reported and can also be detrimental and negatively impact navigation, diving, foraging, communication, and group cohesion (Wright et al., 2007a). Noise has the potential to induce temporary hearing loss also known as temporary threshold shift (TTS), if it is loud or long enough in duration. In general, the higher the sound level and/or longer the duration, the more likely TTS is to occur. If exposure is prolonged or repeated or even as a result of one very loud noise event, the hearing damage can become permanent, also known as a permanent threshold shift (PTS) (Finneran, 2015; Tyack, 2008; Weilgart, 2007). Although progress has been made in understanding the potential adverse effects of noise on marine mammal hearing, much remains unknown on how exposure to high dB levels and impulsive sounds can result in acute or chronic hearing deficits in marine mammal hearing (Finneran, 2015).

Wright et al. (2007a) integrated the available information on noise and stress responses in terrestrial mammals and contended that this can be extrapolated to cetaceans and other marine mammals because stress responses to noise appears to be highly conserved across all the species studied. Thus, for cetaceans as in other mammals, noise is a stressor that may result in “suppression of reproduction (physiologically and behaviorally), accelerated aging and sickness-like symptoms” (Wright et al., 2007b, p. 275). When dolphins cannot remove themselves from prolonged and loud sounds, in addition to physiological stress, damage to overall fitness can also result (Finneran, 2015; Tyack, 2008; Wright et al., 2007a). Changes in the profile of blood parameters have been linked to the effects of noise in cetaceans, altering immune competence in individuals which may have far-ranging consequences for targeted individuals and populations over time (Romano et al., 2004; Thomas, Kastelein, & Awbrey, 1990).

During drive hunts, the repeated and prolonged transmission of high-amplitude sound generated by the metal trumpets and the noise of the boat engines, creates noise at potentially damaging dB levels for these animals and very likely disrupts communication between individuals and groups. The acoustic onslaught experienced by cetaceans during the drive hunts is significant and may negatively impact them in many ways. Importantly, the cumulative multiple stressors resulting from ear damage and the stress responses to noise itself can affect both survivorship and reproduction, thus impacting both the welfare of individuals and health of populations.

Physical impact: handling, restraint and injury

Once sequestered in the cove, the dolphins are sometimes held up to five days before they are selected for slaughter, segregated and taken to sea pen holding facilities where they will eventually be distributed to aquaria, or released. Injury may occur during this selection process. For those individuals selected for slaughter, they are roped around their tails (caudal peduncle), and then towed to the killing area while still alive (see Supplementary Video 1).

Analysis of video of this activity shows that when dolphins are driven close to shore, they exhibit very rapid respiration rates. When forcibly submersed, they lack the capacity to control the duration of their submersion and display profound distress (Butterworth et al., 2013). Diving mammals are highly adapted for submersion, and capable of relatively long intervals between respirations (compared to terrestrial mammals). Diving animals are also capable of storage of oxygen in blood and tissue, and of bradycardia (slowing of the heart rate), adaptations that can extend the dive interval (Castellini, Davis, & Kooyman, 1988; Eckert, Eckert, Ponganis, & Kooyman, 1989; Williams, Haun, & Friedl, 1999). It is our contention that, despite the formidable adaptations to diving of toothed whales, some cetaceans may be drowned or asphyxiated (i.e., die from failure to be able to breathe through

voluntary apnea, as opposed to entry of water into the lungs) during submersion and that many animals are subject to profound respiratory challenges during this process. Drowning or asphyxiation, resulting from forced restraint, would not be permitted in any accepted slaughter process for terrestrial mammals, and this highly aversive mechanism for killing these air-breathing animals has not been formally assessed during discussions on the ethics of small cetacean drive hunts (Figure 3).

Individual dolphins, including calves, are sometimes released so as not to violate species or seasonal quotas. This selection process involves the rough handling of individuals that are often struggling against the nets or exhausted from the chase and herding. Individuals that are selected for marine parks are held in the cove and often removed days after the chase and slaughter has ended. Video footage from documentaries and other images readily available in the media reveal injured dolphins with obvious physical trauma (i.e., bloodied areas on the body) from the handling and restraint associated with selection for marine parks and relocation to the shoreline (Figure 4). Permission to hunt Pacific white-sided dolphins was granted in 2007 and requires special nets (Hiroyuki, 2011). This species is taken primarily for live capture for aquaria, but the capture process—although conducted offshore and outside the confines of the cove—involves the same chase and capture methods, and potential for injury.




Figure 3. (a) A group of Risso's dolphins is corralled behind nets at the conclusion of a drive hunt. Image: DolphinProject.com/Marna Frida Olsen. (b) After chase and confinement in a cove, live Risso's dolphins are tethered and towed. Image: DolphinProject.com/Marna Frida Olsen. (c) While still alive, struggling dolphins are tethered by their tails for towing to the slaughter area (indicated by smaller circle). Other free-swimming dolphins (indicated by larger circle) are in close proximity to those tethered. Image: AtlanticBlue.de



Figure 4. (a) A dolphin is secured in a net after selection for a marine park. Image: Sea Shepherd Conservation Society. (b) An obviously injured dolphin being restrained during selection for a marine park after a drive hunt. Image: Sea Shepherd Conservation Society

The impacts of physical trauma associated with the chase and herding process is also evidenced by individual dolphins that strand and die after the hunts have concluded, often washing ashore days later (see <https://www.youtube.com/watch?v=IVCHdNc13cc>). According to eye-witness accounts, dolphins selected for release rarely leave the cove on their own—rather, they are chased and forced back out to sea by the fishermen (L. Lemieux, personal communication, August 5, 2018). These individuals are severely traumatized and disoriented from the process of the hunts. Pelagic species such as striped dolphins appear to panic in shallow water, throwing themselves onto sharp rocks as they avoid their pursuers in the cove (Supplementary Video 2). Releasing calves or over-quota animals during the hunt may contribute to other hidden or indirect mortality currently not monitored or quantified. Once released, the fate of these individuals injured or traumatized by the chase, handling, and proximity to the slaughter of their pod mates, is unknown.

International guidelines governing handling before slaughter

Recommendations adopted by consensus among the veterinary authorities of the World Organization for Animal Health (OIE), an intergovernmental organization responsible for improving animal health worldwide, state that all animals slaughtered outside slaughterhouses should be managed to ensure that their transport, lairage, restraint and slaughter is carried out without causing undue stress to the animals, including handling in such a way as to avoid harm, distress or injury (Shimshony & Chaudry, 2005; World Organization for Animal Health, 2011). Under the OIE, Japan has an obligation to uphold these principles. These recommendations also include guidelines that would specifically apply to the methods deployed during the drive hunt process, including prohibition against conscious animals being thrown, dragged or dropped; avoidance of excessive pressure that causes struggling or vocalization in animals; excessive shouting or making loud noises that

may agitate the animals; and suspending or hoisting animals by the feet or legs. The handling, restraint and transportation of cetaceans violates these guidelines.

Socio-psychological and ecological impacts

Observational and experimental research conducted in the field and laboratory has demonstrated that dolphins exhibit high levels of cognitive and social complexity, including advanced social and self-awareness, sophisticated social learning, tool use, vocal and social imitation, the ability to comprehend symbolic gestural “sentences”, self-organized learning, metacognition, and cultural transmission (Connor, Watson-Capps, Sherwin, & Krützen, 2011; Herman, Richards, & Wolz, 1984; Herman, 2002; Krützen et al., 2005; Mann & Singh, 2015; Mann & Smuts, 1999; Marino et al, 2007; Reiss & Marino, 2001; Reiss and McCowan, 1993; Smith et al, 1995; Whitehead, Rendell, Osborne, & Wursig, 2004). Notably, advanced social awareness has been well-documented in young dolphins (Connor & Krützen, 2015; Gibson & Mann, 2008; Mann & Smuts, 1999) and young bottlenose dolphins show the emergence of self-awareness, demonstrated by mirror-self recognition, at an age earlier than reported for children and apes (Morrison & Reiss, 2018). Dolphins in Shark Bay, Australia, use sponges as tools during foraging. The practice of “sponging” appears to be transmitted culturally and has been attributed to an almost exclusive vertical social transmission within a single matriline from mother to female offspring (Krützen et al., 2005). Bottlenose dolphins live in complex social groups, exhibiting fission-fusion type social organization in which the size and composition of groups are dynamic, and social factors, such as alloparental care, can contribute to variation in individual reproductive fitness in this species (Frère et al., 2010). Dolphins are mutually-dependent, form long-lasting social alliances that may be more complex than exhibited by other non-human species, and exhibit high levels of social awareness and social cognition (Connor & Krützen, 2015; Connor, 2007). Dolphins are social learners and have an extended period of juvenile dependency during which time youngsters

are cared for by, and learn from, their mothers and other members of the social group (Connor, Read, & Wrangham, 2000; McCowan & Reiss, 1995; Reiss, McCowan, & Marino, 1997; Whitehead, 2010). Although precocious at birth, young dolphins are highly vulnerable, requiring a high degree of parental care and energy investment by mothers and alloparents (Gibson & Mann, 2008; Gubbins, McCowan, Lynn, Hooper, & Reiss, 1999; Noren, 2008).

Dolphins show altruistic and epimeletic (caregiving) behavior towards members of their group and other cetacean species, including assisting pregnant females during delivery, alloparenting, and supporting injured or ill animals (Bearzi et al., 2018; Brown et al., 1966; Cockcroft & Sauer, 1990; Connor & Norris, 1982; Kuczaj et al., 2015; Park et al., 2013; Pilleri, 1984; Wells, 1991; Whitehead, 1996). Dolphins emit distress calls (Lilly, 1963) and the emission of these calls has been reported to elicit epimeletic behavior in other members of the social group (Kuczaj et al., 2015). Vocalizations of distressed dolphins during these hunts can have a psychological and physiological impact on other individuals in the group.

This high level of social awareness in both adult and young dolphins supports the view that the chase, encirclement, capture and removal of individuals from their social groups can have both negative psychological and social impacts. During the hunt process, dolphins are in close proximity to each other when individuals are removed from the group or killed. Individuals held for selection or slaughter are likely to be in a state of extreme anxiety, when they experience and witness the capture and killing of others individuals to whom they may be related or share close social bonds (Butterworth et al., 2013).

The prolonged chasing and capture of dolphins harms not only the individual captured, but can negatively impact the social group and threaten populations and the marine ecosystem (Reeves, Smith, Crespo, & Notarbartolo di Sciara, 2003). The capture of even a few animals may result in the death or injury of many more, since capture involves intensive

harassment of dolphin groups. The removal of individuals that model important behavior and transmit specific information within the social group (e.g., caring for young, demonstrating foraging techniques, or locating critical feeding habitats) may have a negative impact on populations. The transmission of information between generations is beginning to be recognised as non-human culture (Lusseau & Newman, 2004; Wells, 2003; Whitehead et al., 2004; Whitehead, 2011; Whitehead & Rendell, 2015; Williams & Lusseau, 2006).

Noren & Edwards (2007) note the significant potential for mother-calf separation, and eventual calf mortality, resulting from chase and encirclement associated with the tuna purse-seine fishery. In these fisheries, dolphins are often chased, encircled and entangled in nets during the course of fishing operations. During fishery evasion, the potential for unobserved calf mortality is quite high if calves are not reunited promptly with their mother. Beyond direct injury and physical separation, even if young calves and neonates are reunited with their mothers during fishing operations, they may also be affected by an influx of stress hormones delivered through the mothers' milk in response to this intermittent but severe experience. The resulting impact on young dolphins could be amplified due to their own immature nervous systems (Noren & Edwards, 2007). Wright et al. (2007) note that "these excessive stress-chemical loads have the potential for generating both acute neurological damage and long-lasting neurological re-programming in any nursing calves in evasion of a tuna purse-seine set in the ETP." (p. 286).

Thus, drive-type hunts and procedures involving chase and encirclement have a great capacity to disrupt social structures and severely impact social groups and populations (Edwards, 2002a; Edwards 2002b; Scillitani, Monaco, & Toso, 2010). Disruption to social structures may have conservation implications in excess of the number of individuals removed. For example, Frère et al. (2010) examined genetic and social effects on female calving success (a partial measure of fitness) in bottlenose dolphins. They determined that

both genetic and social factors contribute to variation in individual fitness related to female calving success and suggest that the influence of social relationships between females is consistent with either the social transmission of reproductive prowess, or with a type of homophily in which females with calves associate with other females with calves, thus demonstrating the importance of social relationships to sustainability.

Behavioral and physiological impacts

In reviewing available research on behavioral and physiological impacts of chase and hunt activity on both terrestrial and marine mammals, there is a relative lack of literature on the behavioral and physiological impacts of chase, herding and capture of cetaceans when compared to this area of scientific inquiry in other terrestrial species.

Behavioral responses to stressors can be assessed, but only when individual animals can be observed (Eskesen et al., 2009). During the dolphin drives hunters often actively prevent observation by outsiders. Despite this, the acquisition of investigative video has enabled analysis and documented escape and avoidance behaviors which take place during the chase, herding, separation and collection of individuals during the hunt process (Butterworth et al., 2013).

There are multiple influences on the behavioral responses of the driven animals - including vessel speed, the ‘intensity’ of the sound, the water depth and bathymetry of the ocean, the age, sex and ‘experience’ of the animal, size and composition of the social group—all of which influence stress responses (Kastelein et al., 2000; St. Aubin & Dierauf, 2001; Williams et al., 2017). Social instability, changes in respiratory rate, dive patterns, changed posture, and altered vocalizations are short-term responses to stress in cetaceans (Frohoff, 2004; Waples & Gales, 2002). Behavioral changes resulting from stress or ‘distress’ increase vulnerability and risk of predation, and some species may mask such behaviors to

reduce ‘vulnerability’ (Broom, 1991; Esch et al., 2009; Frohoff, 2004; Lusseau, 2003; Pollard et al., 1993; Waples & Gales, 2002).

A stressor can be physical or psychological, and an organism’s response can be acute or chronic. It is understood that mammalian responses have a common neurochemical pathway (Romano et al., 2002; Sapolsky, 1992). When stressed, the hypothalamic-pituitary-adrenal (HPA) axis is stimulated and corticotropin releasing factor (CRF) triggers release of the adreno-corticotropin (ACT) from the pituitary. This in turn induces the secretion of hormones known as adrenocorticoids (glucocorticoids and mineralocorticoids). Although cortisol is the primary glucocorticoid hormone released by this process in most mammals, the presence of corticosterone in adrenocortical tissue has been reported in cetaceans (Carballeira, Brown, Fishman, Trujillo, & Odell, 1987). Glucocorticoids and catecholamines (epinephrine and norepinephrine) cause changes in organ function which include altered rates of gluconeogenesis, and increased heart rate, which contribute to preparedness for ‘fight or flight’ and alterations in the production and release of white blood cells (Guyton & Hall, 2000; Landsberg & Young, 1978; Randall, Burggren, & French, 2002). The effects of cortisol in the animal can be variable, dependent on the context for the animal, including the timing of the stressor, reproductive cyclicity and seasonality (Dobson et al., 2000; Bechshoft et al., 2013), and social and environmental circumstances (Schmitt et al., 2010). The ‘pattern’ of glucocorticoid hormone release may be as important as the amount of hormone released, as the biological effects of the hormones response may take time, and result from hormone receptor interactions which occur across the sometimes-prolonged time duration of the stress response (Laugero, Bell, Bhatnagar, Soriano, & Dallman, 2001; Sheriff, Dantzer, Delehanty, Palme, & Boonstra, 2011).

The effects of glucocorticoids such as cortisol and corticosterone are to generate physiological, behavioral and even anatomical responses to stressors, to affect the

endocrinology of the reproductive system, to control energy metabolism, and to impact growth and development in the young animal. Glucocorticoids have been used as indicators of stress responses for a wide range of mammals (Sheriff et al., 2011; Atkinson et al., 2015) and has been collected and analyzed from a variety of tissue and biological materials including blood, urine, faeces and hair (Macbeth et al., 2012) in terrestrial mammals, and from blood, muscle, blubber (Trana et al., 2015; Kellar et al., 2015), faeces (Palme et al., 2013) and from the liquid- cellular component of the blow (Thompson et al., 2014) in cetaceans. Glucocorticoid analyses, sometimes linked to analysis of behavioral change, have been used to increase understanding of the negative impact of human activity in a number of wild animal species, including: hair cortisol analysis for environmental stressors in deer (Caslini et al., 2016); for assessment of environmental change in polar bears (Weisser et al., 2016); ‘landscape fear’ (human persecution) in brown bears (Stoen et al., 2015); capture and handling in brown bears (Cattet et al., 2014); the effects of human harassment in wolves (Bryan et al., 2015); the effects of poaching on elephants (Gobush et al., 2008); tourist pressures on Chamois (Zwijacz-Kozica et al., 2013); and culling and hunting methods in red deer (Cockram et al., 2011) and in other ungulates (Gentsch et al., 2018).

In the absence of direct studies on the behavioral and stress physiology effects of hunting on cetacean species, it is possible to make inferences of the likely impacts of hunting activity from studies which have looked at non-hunting domains of cetacean-human interactions such as whale watching (Avila et al., 2015; Christiansen et al., 2013), or the impacts of anthropogenic sound (Ellison et al., 2012). In the case of hunted cetaceans, it is possible to make ‘informed inferences’ from studies carried out on terrestrial animals where they are hunted or subject to other stressors of human origin.

In cetaceans, physiological impacts from chase and capture may be acute or chronic. Individuals may become entangled in the capture nets and suffocate or suffer fatal stress-

related conditions associated with the trauma of capture (Curry, 1999; Schmitt, St. Aubin, Schaefer, & Dunn, 2010; St. Aubin & Geraci, 1988; Thompson & Geraci, 1986). Mortality rates of captured bottlenose dolphins has been shown to increase six-fold immediately after capture and do not decrease to 'normal' levels for up to 45 days suggesting an acute physiological response to the event (Small & DeMaster, 1995).

As well as glucocorticoid hormones circulating in blood and tissues, it is possible that stressors result in measurable changes in more structural cellular elements such as complementary DNA (cDNA). Mancina, Warr, and Chapman (2008) describe a cDNA microarray study, in which analysis of sequence tags (ESTs) which were isolated and sequenced, suggested that the stress response and immune function genes of the bottlenose dolphin could be altered by stressors. The immune system of cetaceans can also be compromised by the stress associated with capture and chase, and can lead to premature death (Romano et al., 2002).

Capture myopathy

During drive hunts, small cetaceans swim in ways that subject them to both acute, and more chronic prolonged stressors for which they are not well adapted. 'Capture myopathy' or 'exertional myopathy' (EM) has been described in a number of species (Bateson & Bradshaw, 1997; Chivers & Myrick, 1991; Curry, 1999; Hartup, Kollias, Jacobsen, Valentine, & Kimber, 1999; Jarrett, Jennings, Murray, & Harthoorn, 1964; Maas, 2000; Maas, 2003; Myrick & Perkins, 1994). Exertional myopathy affects skeletal and cardiac muscle, resulting in muscle necrosis (death of muscle fibers), mitochondrial damage, altered polysaccharide storage in the muscle, and destruction of striated skeletal muscle (Aleman, 2008).

Stress-response blood biochemical profiles observed in offshore spotted and spinner dolphins revealed the presence of acute stress-responses in chased and captured dolphins which include cardiac lesions and other tissue damage (Forney, St. Aubin, & Chivers, 2002). During chase and capture of other hunted species (e.g., deer), disruption of muscle tissue, elevated secretion of endorphins, and high concentrations of cortisol are all typically associated with extreme physiological and psychological stress (Bateson & Bradshaw, 1997). Exertional myopathy stemming from the stress and trauma of chase and capture is associated with a release of high levels of creatine kinase, serum aspartate, and other enzymes resulting from the degradation of cardiac and other muscle tissue which can result in compromised health and even death (Broom & Johnson, 1993; Cattet, Stenhouse, & Bollinge, 2008; Fair & Becker, 2000; Maas, 2003; Myrick & Perkins, 1994; St. Aubin, 2002; St. Aubin et al., 2013).

Exertion associated with hunting red deer (*Cervus elaphus*) resulted in severe physiological disturbances in the animals, and many of the changes appeared seriously maladaptive (Bateson & Bradshaw, 1997). The deer, in many cases, had been pushed beyond limits for which they were physically prepared when subjected to prolonged chases. Lesions consistent with exertional myopathy were documented after postmortems of North American river otters (*Lutra canadensis*) which had been caught and then translocated (Hartup et al., 1999). Results showed that 28% of the otters, which were captured and transported over long distances, and then subjected to a brief period of captivity before release, had abnormal elevations in serum enzymes consistent with myopathy. Exertional myopathy has also been described in captured grizzly bears (*Ursus arctos*) following capture using leg-hold snares in Canada (Cattet et al., 2008). In a number of non-cetacean species, capture myopathy has resulted in death after exertion from complicating factors such as heart failure or partial or complete paralysis (Bartsch, McConnell, Imes, & Schmidt, 1977; Beringer, Hansen, Wilding, Fischer, & Sheriff, 1996; Curry, 1999). In cetaceans, capture myopathy (Spraker, 1993),

stress cardiomyopathy (Cebelin & Hirsch, 1980; Colgrove, 1978) and contraction band necrosis, a form of necrosis in cardiac myocytes associated with acute cellular damage (Turnbull & Cowan, 1998), have all been linked to events including capture, transportation, violent assault on cetaceans by humans, and beach stranding in cetaceans.

Many of the scientific findings regarding the physiological impacts associated with chase and encirclement are based on studies conducted as part of a larger research program mandated under the 1997 International Dolphin Conservation Program Act (IDCPA) of the US Marine Mammal Protection Act which have investigated whether the eastern tropical Pacific tuna fishery negatively impacted dolphin stocks. These studies were known collectively as the Chase Encirclement Stress Studies (CHESS) (Forney et al., 2002), and evaluated the degree of acute (i.e., rapid onset, short duration) physiological stress represented by single chase and capture events on cetaceans in the tuna purse seine fishery (Westgate et al., 2007).

The findings of the CHESS and other marine and non-marine mammal studies suggest that chase and encirclement of dolphins results in measurable, and relatively consistent stress responses, in addition to other impacts including physical injury, separation of mothers and calves, and impaired reproductive success (Cramer, Perryman, & Gerrodette, 2008; Edwards, 2006; Matsuda et al., 1996; Noren & Edwards, 2007; St. Aubin et al., 2013). Elevated blood cortisol levels, raised ACTH and catecholamine levels, some elevation of enzymes released from muscle following exertion, and altered white blood cell changes (a shift in the balance of white cell types and an overall net increase in white cell count, the “stress leucogram”) are also reported in the CHESS studies (St. Aubin, 2002).

Conclusion

There is compelling scientific evidence that the process of chase and capture is inhumane. Based on a review of the available scientific data, we contend that the process of drive hunts involving chase, encirclement, and live capture of cetaceans results in physical injury, and physiological and socio-psychological stress which can have enduring negative impacts for individuals and populations. Informed by the CHESS studies, it is our contention that drive hunts, including the chase, subsequent corralling, and handling and selection can have profound negative effects on cetaceans. Based on current knowledge of these animals and the high probability for injury and suffering resulting from pursuit, chase and herding, the authors assert that the drive hunt method cannot be conducted in a humane manner and should be abandoned.

Remarkably, the potential negative impacts of the chase and herding aspects of the drive hunts are absent from policy discussions relating to the methods used to collect dolphins from the wild for marine parks or aquaria. There is a striking disparity between the information we have reviewed in this paper and policymaking decisions regarding the hunting and collection of small cetaceans in general. Noting that advancements in animal welfare policy in modern zoological institutions are science-based, we encourage zoological institutions to acknowledge the adverse impacts of chase and capture on individuals and populations of delphinid species during acquisition from the wild. We argue that based upon the evidence provided herein, the live collection of any cetaceans from the wild should be discontinued, or at least reconsidered.

Supplementary videos (S1 and S2)

Contact first or corresponding author for videos at: courtney@lightkeepersfoundation.com, or Diana Reiss, dlr28@columbia.edu

References

- Akamatsu, T., Hatakeyama, Y., & Takatsu, N. (1993). Effects of pulse sounds on escape behaviour of false killer whales. *Nippon Suisan Gakkaishi, Bulletin of the Japanese Society of Scientific Fisheries*, 59(8), 1297-303.
- Aleman, M. (2008). A review of equine muscle disorders. *Neuromuscular Disorders*, 18, 277-287.
- Atkinson, S., Crocker, D., Houser, D., & Mashburn, K. (2015). Stress physiology in marine mammals: how well do they fit the terrestrial model? *Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology*, 185, 463–486.
- Avila, I. C., Correa, L. M., & Parsons, E. C. M. (2015). Whale-watching activity in Bahía Málaga, on the Pacific Coast of Colombia, and its effect on humpback whale (*Megaptera novaeangliae*) behavior. *Tourism in Marine Environments*, 11, 19–32.
- Bartsch, R. C., McConnell, E. E., Imes, G. D., & Schmidt, J. M. (1977). A review of exertional rhabdomyolysis in wild and domestic animals and man. *Veterinary Pathology*, 14, 314-0324.
- Bateson, P., & Bradshaw E. L. (1997). Physiological effects of hunting red deer (*Cervus elaphus*). *Proceedings of the Royal Society of London B*, 264, 1707-1714.
- Bearzi, G., Kerem, D., Furey, N. B., Pitman, R. L., Rendell, L., & Reeves, R.R. (2018). Whale and dolphin behavioural responses to dead conspecifics. *Zoology*, 128, 1-15.
- Bechshoft, T.O., Sonne, C., Riget, F.F., Letcher, R.J., Novak, M.A., Henschey, E., Meyer, J.S., Eulaers, I., Jaspers, V.L.B., Covaci, A., & Dietz, R. (2013). Polar bear stress hormone cortisol fluctuates with the North Atlantic Oscillation climate index. *Polar Biology*, 36, 1525–1529.

- Beringer, J., Hansen, L. P., Wilding, W., Fischer, J., & Sheriff, S. L. (1996). Factors affecting capture myopathy in white-tailed deer. *Journal of Wildlife Management*, 60, 373-380.
- Broom, D. M. (1991). Animal welfare: Concepts and measurements. *Journal of Animal Science*, 69, 4167-4175.
- Broom, D. M., & Johnson, K. G. (1993). Stress and animal welfare. London: Chapman & Hall.
- Brown, D. H., Caldwell, D. K., & Caldwell, M. C. (1966). Observations on the behavior of wild and captive false killer whales with notes on associated behavior of other genera of captive delphinids. *Los Angeles County Museum Contributions in Science*, 95, 1-32.
- Brownell, R. L., Nowacek, D. P., & Ralls, K. (2008). Hunting cetaceans with sound: a worldwide review *Journal of Cetacean Research and Management*, 10(1), 81-88.
- Bryan, H. M., Smits, J. E. G., Koren, L., Paquet, P., Wynne-Edwards, K. E., & Musiani. (2015). Wolves that are heavily hunted have higher stress and reproductive hormones than wolves with lower hunting pressure. *Functional Ecology*, 29(3), 347-356
- Butterworth, A., Brakes, P., Vail, C. S., & Reiss, D. (2013). A veterinary and behavioral analysis of dolphin killing methods currently used in the "drive hunt" in Taiji, Japan. *Journal of Applied Animal Welfare Science*, 16(2), 184-204.
- Carballeira, A., Brown, J. W., Fishman, L. M., Trujillo, D., & Odell, D. K. (1987). The adrenal gland of stranded whales (*Kogia breviceps* and *Mesoplodon europaeus*): Morphology, hormonal contents, and biosynthesis of corticosteroids. *General and Comparative Endocrinology*, 68(2), 293-303.

- Caslini, C., Comin, A., Peric, T., Prandi, A., Pedrotti, L., & Mattiello, S. (2016). Use of hair cortisol analysis for comparing population status in wild red deer (*Cervus elaphus*) living in areas with different characteristics. *European Journal of Wildlife Research*, 62, 713.
- Castellini, M. A., Davis, R. W., & Kooyman, G. L. (1988). Blood chemistry regulation during repetitive diving in Weddell seals. *Physiological and Biochemical Zoology*, 61(5), 379-386.
- Cattet, M., Stenhouse, G., & Bollinge, T. (2008). Exertional myopathy in a Grizzly Bear (*Ursus arctos*) captured by leghold snare. *Journal of Wildlife Diseases*, 44(4), 973–978.
- Cattet, M., Macbeth, B. J., Janz, D. M., Zedrosser, A., Swenson, J. E., Dumond, M., & Stenhouse, G. B. (2014). Quantifying long-term stress in brown bears with the hair cortisol concentration: A biomarker that may be confounded by rapid changes in response to capture and handling. *Conservation Physiology*, 2, 1–15.
- Cebelin, M. S., & Hirsch, C. S. (1980). Human stress cardiomyopathy. *Human Pathology*, 11(2), 123-132.
- Chivers, S. J., & Myrick, A. C. (1991). Comparison of age at sexual maturity and other reproductive parameters for two stocks of spotted dolphins, *Stenella attenuata*. *Fishery Bulletin*, 91, 611-618.
- Christiansen, F., Rasmussen, M. H., & Lusseau, D. (2013). Whale watching disrupts feeding activities of minke whales on a feeding ground. *Marine Ecology Progress Series*, 478, 239-251.
- Cockcroft, V. G., & Sauer, W. (1990). Observed and inferred epimeletic (nurturant) behavior in bottlenose dolphins. *Aquatic Mammals*, 16.1, 31-32.

- Cockram, M. S., Shaw, D. J., Milne, E., Bryce, R., McClean, C., & Daniels, M. J. (2011). Comparison of effects of different methods of culling red deer (*Cervus elaphus*) by shooting on behaviour and post mortem measurements of blood chemistry, muscle glycogen and carcass characteristics. *Animal Welfare*, 20(2), 211–224.
- Colgrove, G. S., (1978). Suspected transportation-associated myopathy in a dolphin. *Journal of the American Veterinary Medical Association*, 173(9), 1121-1123.
- Connor, R. C., & Norris, K. S. (1982). Are dolphins reciprocal altruists? *The American Naturalist*, 119, 358-374.
- Connor, R. C., Read, A. J., & Wrangham, R. (2000). Male reproductive strategies and social bonds. In J. Mann, R.C. Connor, P.J. Tyack, & H. Whitehead (Eds.), *Cetacean Societies: Field Studies of Dolphins and Whales* (pp. 247-269). Chicago, Illinois: University of Chicago Press.
- Connor, R. C., Watson-Capps, J. J., Sherwin, W. B., & Krützen, M. (2011). A new level of complexity in the male alliance networks of Indian Ocean bottlenose dolphins (*Tursiops sp.*). *Biology Letters*, 7(4), 623-626.
- Connor, R.C., & Krützen, M. (2015). Male dolphin alliances in Shark Bay: Changing perspectives in a 30-year study. *Animal Behaviour*, 103, 223-235. DOI: 10.1016/j.anbehav.2015.02.019
- Cramer, K. L., Perryman, W. L., & Gerrodette, T. (2008). Declines in reproductive output in two dolphin populations depleted by the yellowfin tuna purse-seine fishery. *Marine Ecology Progress Series*, 369, 273-285.
- Curry, B. (1999). Stress in mammals: The potential influence of fishery-induced stress on dolphins in the eastern tropical Pacific Ocean. Report to Congress, U.S. Department

of Commerce NOAA Technical Memo 260, pp. 1-56.

Dobson, H., & Smith, R.F. (2000). What is stress, and how does it affect reproduction?

Animal Reproductive Science, 60, 743–752.

Eckert, S. A., Eckert, K. L., Ponganis, P., & Kooyman, G. L. (1989). Diving and foraging behavior of leatherback sea turtles (*Dermochelys coriacea*). *Canadian Journal of Zoology*, 67, 2834-2840.

Edwards, E. F. (2002a). Energetics consequences of chase by tuna purse-seiners for spotted dolphins (*Stenella attenuata*) in the Eastern Tropical Pacific Ocean. Southwest Fisheries Science Center, La Jolla, CA 92037.

Edwards, E. F. (2002b). Behavioral contributions to separation and subsequent mortality of dolphin calves chased by tuna purse-seiners in the eastern tropical Pacific Ocean. Southwest Fisheries Science Center. La Jolla, CA 92037

Edwards, E. F. (2006). Duration of unassisted swimming activity for spotted dolphin (*Stenella attenuata*) calves: implications for mother-calf separation during tuna purse-seine sets. *Fishery Bulletin*, 104, 125–135.

Ellison, W. T., Southall, B. L., Clark, C. W., & Frankel, A. S. (2012). A new context-based approach to assess marine mammal behavioral responses to anthropogenic sounds. *Conservation Biology*, 26, 21–28.

Esch, C. H., Sayigh, L. S., Blum, J. E., & Wells, R. S. (2009). Whistles as potential indicators of stress in bottlenose dolphins (*Tursiops truncatus*). *Journal of Mammalogy*, 90(3), 638-650.

Eskesen, I. G., Teilmann, J., Geertsen, B. M., Desportes, G., Riget, F., Dietz, R., & Larsen, L. (2009). Stress level in wild harbour porpoises (*Phocoena phocoena*) during satellite tagging measured by respiration, heart rate and cortisol. *Journal of the Marine Biological Association of the United Kingdom*. 89(5), 885–892.

doi:10.1017/S0025315408003159

- Fair, P. A., & Becker, P. R. (2000). Review of stress in marine mammals. *Journal of Aquatic Ecosystem Stress and Recovery*, 7, 335-354.
- Finneran, J. J. (2015). Noise-induced hearing loss in marine mammals: A review of temporary threshold shift studies from 1996 to 2015. *The Journal of the Acoustical Society of America*, 138, 1702.
- Forney, K. A., St. Aubin, D. J., & Chivers, S. J. (2002). Chase encirclement stress studies on Dolphins involved in eastern tropical Pacific Ocean purse seine operations during 2001 (Southwest Fisheries Science Center Administrative Report). La Jolla, CA: National Marine Fisheries Service.
- Frère, C. H., Krützen, M., Mann, J., Connor, R. C., Bejder, L., & Sherwin, W.B. (2010). Social and genetic interactions drive fitness variation in a free-living dolphin population. *Proceedings of the National Academy of Sciences of the United States of America*, 107(46), 19949–19954.
- Frohoff, T. G. (2004). Stress in dolphins. In M. Beckoff (Ed.) *Encyclopedia of animal behavior* (pp. 1158-1164). Westport, CT: Greenwood Press.
- Gentsch, R. P., Kjellander, P., & Röken, B. O. (2018). Cortisol response of wild ungulates to trauma situations: Hunting is not necessarily the worst stressor. *European Journal of Wildlife Research*, 64, 11.
- Gibson, Q. A., & Mann, J. (2008). Early social development in wild bottlenose dolphins: sex differences, individual variation and maternal influence. *Journal of Animal Behaviour*, 76, 375-387.
- Gobush, K. S., Mutayoba, B. M., & Wasser, S. K. (2008). Long term impacts of poaching on relatedness, stress physiology, and reproductive output of adult female African elephants. *Conservation Biology*, 22, 1590–1599.

- Guyton, A., & Hall, J. (2000). Energetics and metabolic rate. In A.C. Guyton, & J. E. Hall (Eds.), *Textbook of Medical Physiology* (pp. 859-863). Philadelphia, PA: Saunders.
- Hartup, B. K., Kollias, G. V., Jacobsen, M. C., Valentine, B. A., & Kimber, K. R. (1999). Exertional Myopathy in Translocated River Otters from New York. *Journal of Wildlife Diseases*, 35(3), 542-547.
- Herman, L. M., Richards, D. G., Wolz, J. P. (1984). Comprehension of sentences by bottlenosed dolphins. *Cognition*, 16, 129–219.
- Herman, L. H. (2002). Vocal, social, and self-imitation by bottlenosed dolphins. In K. Dautenhahn & C. L. Nehaniv (Eds.), *Complex adaptive systems. Imitation in animals and artifacts* (pp. 63-108). Cambridge, MA: MIT Press.
- Hiroyuki, M. (2011). A case study of small scale coastal whaling of actual use, Wakayama Prefecture Taiji (Whales: resources for transformation). In *Anthropology of the marine environment conservation*. National Museum of Ethnology Survey Report 97: 237Q267, Endo Aiko Ocean Policy Research Foundation policy research group.
- Jarrett, W. F. H., Jennings, F. W., Murray, M., & Harthoorn, A. M. (1964). Muscular dystrophy in a wild Hunter's antelope. *East African Wildlife Journal*, 2, 158-159.
- Kasuya, T. (2002). Japanese whaling. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (pp. 655-662). San Diego, CA: Academic Press.
- Kasuya, T. (2007) Japanese whaling and other cetacean fisheries. *Environmental Science and Pollution Research*, 14(1), 39-48.

- Kellar, N. M., Catelani, K. N., Robbins, M. N., Trego, M. L., Allen, C. D., Danil, K., & Chivers, S. J. (2015). Blubber cortisol: a potential tool for assessing stress response in free-ranging dolphins without effects due to sampling. *PLoS ONE*, *10*, e0115257. doi: 10.1371/journal.pone.0115257. PMID: 25643144.
- Ketten, D. R. (2016). Marine Mammal Auditory System Noise Impacts: Evidence and Incidence. In A.N. Popper & A. Hawkins (Eds.), *The Effects of Noise on Aquatic Life. Advances in Experimental Medicine and Biology* (pp. 207-212). New York, NY: Springer.
- Krützen, M., Mann, J., Heithaus, M. R., Connor, R. C., Bejder, L., Sherwin, W. B. (2005). Cultural transmission of tool use in bottlenose dolphins. *Proceedings of the National Academy of Sciences*, *102* (25), 8939-8943.
- Kuczaj, S. A., Frick, E. E., Jones, B. L., Lea, J. S. E; Beecham, D., & Schnoller, F. (2015). Underwater observations of dolphin reactions to a distressed conspecific. *Learning & Behavior*, *43*(3), 289-300.
- Landsberg, L., & Young, J. B. (1978). Fasting, feeding and regulation of the sympathetic nervous system. *The New England Journal of Medicine*, *298*, 1295-1301.
- Laugero, K. D., Bell, M. E., Bhatnagar, L., Soriano, M., & Dallman, M. F. (2001). Sucrose ingestion normalizes central expression of corticotropin-releasing-factor messenger ribonucleic acid and energy balance in adrenalectomized rats: A glucocorticoid metabolic-brain axis? *Endocrinology*, *142*(7), 2796-2804.
- Lilly, J. C. (1963). Distress call of the bottlenose dolphin: stimuli and evoked behavior responses. *Science*, *139*(3550), 116-8.

- Lusseau, D. (2003). Effects of tour boats on the behaviour of bottlenose dolphins: Using Markov chains to model anthropogenic impacts. *Conservation Biology*, 17(6), 1785–1793.
- Lusseau, D., & Newman, M. E. J. (2004). Identifying the role that animals play in their social networks. *Proceedings of the Royal Society B: Biological Sciences, Biology Letters* (Supplement 6), 271, S477-S481.
- Maas, B. (2000). Prepared and shipped: A multidisciplinary review of the effects of capture, handling, housing and transport on morbidity and mortality. A Report for the RSPCA, 55pp.
- Maas, B. (2003). The potential stress effects of whaling operations and the welfare implications for hunted cetaceans. IWC/55/WK19.
- Macbeth, B. J., Cattet, M. R. L., Obbard, M. E., Middel, K., & Janz, D. M. (2012). Evaluation of hair cortisol concentration as a biomarker of long-term stress in free ranging polar bears. *Wildlife Society Bulletin*, 36, 747–758.
- Mancia, A., Warr, G. W., & Chapman, R. W. (2008). A transcriptomic analysis of the stress induced by capture–release health assessment studies in wild dolphins (*Tursiops truncatus*). *Molecular Ecology*, 17, 2581–2589.
- Mann, J., & Singh, L.O. (2015). Culture, diffusion, and networks in social animals. In R. Scott, & S. Kosslyn (Eds.), *Emerging Trends in the Social and Behavioral Sciences* (pp. 1-16). Hoboken, NJ: John Wiley & Sons.
- Mann, J., & Smuts, B. (1999). Behavioral development in wild bottlenose dolphin newborns (*Tursiops sp.*). *Behaviour*, 136, 529-566.
- Marino, L. M., Connor, R. C., Fordyce, E., Herman, L. M., Hof, P. R., Lefebvre, L., Lusseau, D., McCowan, B., Nimchinsky, E. A., Pack, A. A., Rendell, L., Reidenberg, J. S.,

- Reiss, D., Uhen, M. D., Van der Gucht, E., & Whitehead, H. (2007). Cetaceans have complex brains for complex cognition. *PLoS Biology*, 5(5), 996-991.
- Matsuda, D. M., Bellem, A. C., Gartley, C. J., Madison, V., King, W. A., Liptrap, R. M. & Goodrowe, K. L. (1996). Endocrine and behavioral events of estrous cyclicity and synchronization in wood bison (*Bison bison athabasca*). *Theriogenology*, 45(8), 1429-1441.
- McCowan, B., & Reiss, D. (1995). Whistle contour development in captive-born infant bottlenose dolphins: Role of learning. *Journal of Comparative Psychology*, 109(3), 242-260.
- Morrison, R., & Reiss, D. (2018). Precocious development of self-awareness in dolphins. *PLoS ONE*, 13(1), e0189813. Available at <https://doi.org/10.1371/journal.pone.0189813>
- Myrick, A. C. & Perkins, P. C. (1994). Adrenocortical color and correlates as indicators of continuous acute stress before death in force-chased and purse-seined dolphins. *Pathophysiology*, 1, 390.
- Noren, S. R., & Edwards, E. F. (2007). Physiological and behavioral development in delphinid calves: Implications for calf separation and mortality due to tuna purse seine nets. *Marine Mammal Science*, 23(1), 15-29.
- Noren, S. R. (2008). Infant carrying behaviour in dolphins: costly parental care in an aquatic environment. *Functional Ecology*, 22(2), 284–288.
- Nowacek, D. P., Thorne, L. H., Johnston, D. W., & Tyack, P. L. (2007). Responses of cetaceans to anthropogenic noise. *Mammal Review*, 37, 81–115. doi:10.1111/j.1365-2907.2007.00104.x

- Ohsumi, S. (1972). Catch of marine mammals, mainly of small cetaceans by local fisheries along the coast of Japan. *Bulletin Far Seas Fisheries Research Laboratory*, 7, 137-66.
- Palme, R., Touma, C., Arias, N., Dominchin, M. F., & Lepschy, M. (2013). Steroid extraction: Get the best out of faecal samples. *Wiener Tierärztliche Monatsschrift*, 100, 238–246.
- Park, K. J., Sohn, H., An, Y. R., Moon, D.Y., Choi, S.G., & An, D. H. (2013). An unusual case of care-giving behavior in wild long-beaked common dolphins (*Delphinus capensis*) in the East Sea. *Marine Mammal Science*, 29, 508–514.
- Pilleri, G. (1984). Epimeletic behavior in cetacea: Intelligent or instinctive. *Investigations on Cetacea*, 16, 30-48.
- Pollard, J. C., Littlejohn, R. P., & Suttie, J. M. (1993). Effects of isolation and mixing of social groups on heart rate and behaviour of red deer stags. *Applied Animal Behavioral Science*, 38, 311–322.
- Randall, D., Burggren, W., & French, K. (2002). *Physiology: Mechanisms and Adaptations*. New York: W. H. Freeman and Company.
- Reeves, R. R., Smith, B. D., Crespo, E. A., & Notarbartolo di Sciara, G. (compilers). (2003). *Dolphins, Whales and Porpoises: 2002-2010 Conservation Action Plan for the World's Cetaceans*. Gland, Switzerland and Cambridge, UK: IUCN/SSC Cetacean Specialist Group. Available at <http://app.iucn.org/dbtw-wpd/edocs/2003-009.pdf>
- Reiss, D., & Marino, L. (2001). Mirror self-recognition in the bottlenose dolphin: A case of cognitive convergence. *Proceedings of the National Academy of Sciences*, 98(10), 5937-5942.

- Reiss, D. and McCowan, B. (1993). Spontaneous vocal mimicry and production by bottlenose dolphins (*Tursiops truncatus*): Evidence for vocal learning. *Journal of Comparative Psychology* 107(3), 301-312.
- Richardson, W. J., & Wursig, B. (1997). Influences of man-made noise and other human actions on cetacean behaviour. *Marine and Freshwater Behavior and Physiology*, 29(1-4), 183-209.
- Romano, T. A., Felten, D. L., Stevens, S. Y., Olschowka, J. A., Quaranta, V., & Ridgway, S. (2002). Immune response, stress, and environment: Implications for cetaceans. In C. J. Pfeiffer (Ed.), *Molecular and cell biology of marine mammals* (pp. 253-279), Malabar, Florida: Krieger Publishing Company.
- Romano, T. A., Keogh, M. J., Kelly, C., Feng, P., Berk, L., Schlundt, C. E., Carder, D. A., & Finneran, J. J. (2004). Anthropogenic sound and marine mammal health: measures of the nervous and immune systems before and after intense sound exposure. *Canadian Journal of Fisheries and Aquatic Sciences*, 61, 1123-1134.
- Sapolsky, R. M. (1992). Stress, the aging brain, and the mechanisms of neuron death. Cambridge, MA: The MIT Press.
- Schmitt, T. L., St. Aubin, D. J., Schaefer, A. M., & Dunn, J. L. (2010). Baseline, diurnal variations, and stress-induced changes of stress hormones in three captive beluga whales, *Delphinapterus leucas*. *Marine Mammal Science*, 26(3), 635-647.
- Scillitani, L., Monaco, A., & Toso, S. (2010). Do intensive drive hunts affect wild boar (*Sus scrofa*) spatial behaviour in Italy? Some evidences and management implications. *European Journal of Wildlife Research*, 56(3), 307-318.

- Sheriff, M. J., Dantzer, B., Delehanty, B., Palme, R., & Boonstra, R. (2011). Measuring stress in wildlife: techniques for quantifying glucocorticoids. *Oecologia*, 166(4), 869-887.
- Shimshony, A., & Chaudry, M. M. (2005). Slaughter of animals for human consumption. *Scientific and Technical Review of the Office International des Epizooties*, 24(2), 693-710.
- Small, R., & DeMaster, D. P. (1995). Acclimation to captivity: a quantitative estimate based on survival of bottlenose dolphins and California sea lions. *Marine Mammal Science*, 11, 510-519.
- Smith, J. D., Schull, J., Strote, J., McGee K., Egnor, R. & Erb, L. (1995) The uncertain response in the bottlenose dolphin (*Tursiops truncatus*). *Journal of Experimental Psychology: General*, 124, 391-408.
- Spraker, T. R. (1993). Stress and capture myopathy in artiodactyls. In M. E. Fowler (Ed.), *Zoo and Wild Animal Medicine* (pp. 481-488). Philadelphia, PA: W. B. Saunders.
- St. Aubin, D. J., & Geraci, J. R. (1988). Capture and handling stress suppresses circulating levels of thyroxine (T4) and Triiodothyronine (T3) in beluga whales, *Delphinapterus leucas*. *Physiological Zoology*, 61, 170-175.
- St. Aubin, D. J., Forney, K. A., Chivers, S. J., Scott, M. D., Danil, K., Romano, T. A., Wells, R. S., & Gulland, F. M. D. (2013). Hematological, serum chemistry, and plasma chemical constituents in pantropical spotted dolphins (*Stenella attenuata*) following chase, encirclement, and tagging. *Marine Mammal Science*, 29(1), 14-35.
- St. Aubin, D. J., & Dierauf, L. A. (2001). *Stress in marine mammals*. In L.A. Dierauf, F. M., & D. Gulland (Eds.), *CRC Handbook of Marine Mammal Medicine* (pp. 253-270). Boca Raton, Florida: CRC Press.

- St. Aubin, D. J. (2002). Hematological and serum chemical constituents in eastern spotted dolphins (*Stenella attenuata*) following chase and encirclement. *Southwest Fisheries Science Center*, La Jolla, CA 92037.
- Stoen, O., Ordiz, A., Evans, Alina, L., Laske, T. G., Kindberg, J., Frobert, O., Swenson, J. E., & Arnemo, J. M. (2015) Physiological evidence for a human induced landscape of fear in brown bears (*Ursus arctos*). *Physiology & Behavior*, 152, 244-248.
- Thomas, J. A., Kastelein, R. A., & Awbrey, F. T. (1990). Behavior and blood catecholamines of captive belugas during playbacks of noise from an oil drilling platform. *Zoo Biology*, 9, 393-402.
- Thompson, C. A., & Geraci, J. R. (1986). Cortisol, aldosterone, and leucocytes in the stress response of bottlenose dolphins, *Tursiops truncatus*. *Canadian Journal of Fisheries and Aquatic Science*, 43, 1010-1016.
- Thompson, L. A., Spoon, T. R., Goertz, C. E. C., Hobbs, R. C., & Romano, T. A. (2014). Blow collection as a non-invasive method for measuring cortisol in the beluga (*Delphinapterus leucas*). *PLoS ONE*, 9, 22.
- Trana, M. R., Roth, J. D., Tomy, G. T., Anderson, W. G., & Ferguson, S. H. (2015). Influence of sample degradation and tissue depth on blubber cortisol in beluga whales. *Journal of Experimental Marine Biology and Ecology*, 462, 8–13.
- Turnbull, B. S., & Cowan, D. F. (1988). Myocardial contraction band necrosis in stranded cetaceans. *Journal of Comparative Pathology*, 118, 317-327.
- Tyack, P. L. (2008). Implications for marine mammals of large-scale changes in the marine acoustic environment. *Journal of Mammalogy*, 89 (3), 549-558.

- Tyack, P.L., and Clark, C.W. (2000). Communication and acoustic behavior of dolphins and whales. In W. Au, A.N. Popper, & R. Fay (Eds.), *Hearing by whales and dolphins* (pp. 156-224). New York: Springer Handbook of Auditory Research Series, Springer Verlag.
- Waples, K. A., & Gales, N. J. (2002). Evaluating and minimizing social stress in the care of captive bottlenose dolphins (*Tursiops truncatus*). *Zoobiology*, 21(1), 5-26.
- Weilgart, L. S. (2007). A Brief Review of Known Effects of Noise on Marine Mammals. *International Journal of Comparative Psychology*, 20, 159-168.
- Weisser, J. J., Hansen, M., Björklund, E., Sonne, C., Dietz, R., & Styrishave, B. (2016). A novel method for analysing key corticosteroids in polar bear (*Ursus maritimus*) hair using liquid chromatography tandem mass spectrometry. *Journal of Chromatography B*, 1017–1018, 45–51.
- Wells, R. S. (1991). Bringing up baby. *Natural History*, August, 56-62.
- Wells, R. S. (2003). Dolphin social complexity: Lessons from long-term study and life history. In F. B. M. de Waal, & P. L Tyack (Eds.), *Animal Social Complexity: Intelligence, Culture, and Individualized Societies* (pp. 32-56). Cambridge, MA: Harvard University Press,
- Westgate, A. G., McLellan, W. A., Wells, R. S., Scott, M. D., Meagher, E. M., & Pabst, D. A. (2007). A new device to remotely measure heat flux and skin temperature from free-swimming dolphins. *Journal of Experimental Marine Biology and Ecology*, 346, 45-59.
- Whitehead, H. (1996). Babysitting, dive synchrony, and indications of alloparental care in sperm whales. *Behavioral Ecology and Sociobiology*, 38, 237-244.

- Whitehead, H., Rendell, L., Osborne, R. W., & Würsig, B. (2004). Culture and conservation of non-humans with reference to whales and dolphins: *Review and new directions in Biological Conservation*, 120, 431-441.
- Whitehead, H. (2011). *The cultures of whales and dolphins*. In P. Brakes & M. P. Simmonds (Eds.), *Whales and Dolphins – Cognition, Culture, Conservation and Human Perceptions* (pp. 149-165). London and Washington: Earthscan.
- Whitehead, H., & Rendell, L. (2015). *The Cultural Lives of Whales and Dolphins*. Chicago, Illinois: University of Chicago Press.
- Williams, R., & Lusseau, D. (2006). A killer whale social network is vulnerable to targeted removals. *Biology Letters*, 2, 497–500.
- Williams, T. M., Haun, J. E., & Friedl, W. A. (1999). The diving physiology of bottlenose dolphins (*Tursiops truncatus*). *The Journal of Experimental Biology*, 202, 2739-2748.
- Williams, T. M., Kendall, T. L., Richter, B. P., Ribeiro-French, C. R., John, J. S., Odell, K. L., Losch, B. A., Feuerbach, D. A., & Stamper, A. M. (2017). Swimming and diving energetics in dolphins: A stroke-by-stroke analysis for predicting the cost of flight responses in wild odontocetes. *Journal of Experimental Biology*, 220, 1135-1145. doi: 10.1242/jeb.154245.
- World Organization for Animal Health. (2011). OIE Terrestrial Animal Health Code, Volume 1, General Provisions. Available at <https://www.oie.int/doc/ged/D10905.PDF>
- Wright, A. J., Aguilar Soto, N, Baldwin, A. L., Bateson, M., Beale, C. M., Clark, C., Deak, T., Edwards, E. F., Fernández, A., Godinho, A., Hatch, L. T., Kakuschke, A., Lusseau, D., Martineau, D., Romero, L. M., Weilgart, L. S., Wintle, B. A., Notarbartolo di Sciara, G., & Martin, V. (2007a). Anthropogenic noise as a stressor in

animals: A multidisciplinary perspective. *International Journal of Comparative Psychology*, 20, 250-273.

Wright, A. J., Aguilar Soto, N., Baldwin, A. L., Bateson, M., Beale, C. M., Clark, C., Deak, T., Edwards, E. F., Fernandez, A., Godinho, A., Hatch, L. T., Kakuschke, A., Lusseau, D., Martineau, D., Romero, L. M., Weilgart, L. S., Wintle, B. A., Notarbartolo-di Sciara, G., & Martin, V. (2007b). Do marine mammals experience stress related to anthropogenic noise? *International Journal of Comparative Psychology*, 20, 274-316.

Zwijacz-Kozica, T., Selva, N., Barja, I., Silván, G., Martínez-Fernández, L., Illera, J. C., & Jodłowski, M. (2013). Concentration of fecal cortisol metabolites in chamois in relation to tourist pressure in Tatra National Park (South Poland). *Acta Theriol*, 58, 215–222.

Figure Captions

Figure 1. A fleet of boats returns from a dolphin drive hunt in Taiji, Japan. Image: DolphinProject.com

Figure 2. a) Metal trumpets utilized in the dolphin drive hunts. b) Trumpets (indicated by circle) are mounted on vessels and placed in the water. Using a hammer or other object, fishermen bang on the poles to create loud noise (up to 205dB) that disorients the dolphins. Image: Courtney Vail

Figure 3. a) A group of Risso's dolphins is corralled behind nets at the conclusion of a drive hunt. Image: DolphinProject.com/Marna Frida Olsen. b) After chase and confinement in a cove, live Risso's dolphins are tethered and towed. Image: DolphinProject.com/Marna Frida Olsen. c) While still alive, struggling dolphins are tethered by their tails for towing to the slaughter area (indicated by smaller circle). Other free-swimming dolphins (indicated by larger circle) in close proximity to those tethered. Image: AtlanticBlue.de

Figure 4. a) A dolphin is secured in a net after selection for a marine park. Image: Sea Shepherd Conservation Society. b) An obviously injured bottlenose dolphin being restrained during selection for a marine park after a drive hunt. Image: Sea Shepherd Conservation Society

Supplemental Video Material

Supplementary Video 1

<https://www.dropbox.com/s/m33kw01r4uy5ft9/S1%20Video%20Tail%20tether.mp4?dl=0>

Supplementary Video 2

<https://www.dropbox.com/s/siime5c85f13fuu/S2%20Video%20Blood%20in%20Taiji%20Cove%2C%20January%2018%2C%202011.mp4?dl=0>